

L613D005/04

FRONT

150mm

13mm

13mm

8mm

5mm

100mm

5mm

25mm

5mm

1st bar

20mm

Last bar

Solpadeine PLUS CAPSULES
Paracetamol, Codeine, Caffeine

This medicine is for the short term treatment of acute moderate pain when other painkillers have not worked.

You should only take this product for a maximum of 3 days at a time. If you need to take it for longer than 3 days you should see your doctor or pharmacist for advice.

This medicine contains codeine which can cause addiction if you take it continuously for more than 3 days. This can give you withdrawal symptoms from the medicine when you stop taking it. If you take this medicine for headaches for more than 3 days it can make them worse.

Please read right through this leaflet before you start using this medicine. This medicine is available without prescription, but you still need to use Solpadeine Plus Capsules carefully to get the best results from them.

Keep this leaflet you may need to read it again.

If you have any questions, or if there is anything you do not understand, ask your pharmacist.

In this leaflet:

1. What Solpadeine Plus Capsules do
2. Check before you take Solpadeine Plus Capsules
3. How to take Solpadeine Plus Capsules
4. Possible side effects
5. How to store Solpadeine Plus Capsules
6. Further information

1. What Solpadeine Plus Capsules do

Solpadeine Plus Capsules can be used in patients over 12 years of age for the short-term relief of moderate pain that is not relieved by other painkillers such as paracetamol or ibuprofen alone.

They can be used for migraine, headache, dental pain, period pain, strains & sprains, backache, rheumatic pain and sciatica. The capsules contain three active ingredients; paracetamol, codeine and caffeine. Codeine and paracetamol are both painkillers. Codeine belongs to a group of medicines called opioid analgesics which act to relieve pain.

It can be used on its own or in combination with other painkillers. Caffeine acts to further help the effectiveness of paracetamol.

2. Check before you take Solpadeine Plus Capsules

- This medicine contains codeine which can cause addiction if you take it continuously for more than 3 days. This can give you withdrawal symptoms from the medicine when you stop taking it.
- If you take a painkiller for headaches for more than 3 days it can make them worse.

continued... 1100000010692

BACK

150mm

13mm

13mm

8mm

5mm

100mm

5mm

25mm

5mm

1st bar

20mm

Last bar

Use in children with breathing problems: Codeine is not recommended in children with breathing problems, since the symptoms of morphine toxicity may be worse in these children.

! Pregnancy and breast feeding

Talk to your doctor before taking Solpadeine Plus Capsules if you are pregnant. Do not take Solpadeine Plus Capsules while you are breast-feeding. Codeine and morphine passes into breast milk.

3. How to take Solpadeine Plus Capsules

Adults and children aged 12 years and over:
Swallow 2 capsules every 4 hours as needed.

! Do not take more frequently than every 4 hours

- Do not take more than 8 capsules in 24 hours
- Do not take more than the recommended dose
- Avoid too much caffeine in drinks like coffee and tea. High caffeine intake can cause difficulty sleeping, shaking and an uncomfortable feeling in the chest
- Solpadeine Plus Capsules should not be taken by children below the age of 12 years, due to the risk of severe breathing problems
- This medicine should not be taken for more than 3 days. If the pain does not improve after 3 days, talk to your doctor for advice.

Possible withdrawal effect

This medicine contains codeine and can cause addiction if you take it continuously for more than 3 days. When you stop taking it you may get withdrawal symptoms. You should talk to your doctor or pharmacist if you think you are suffering from withdrawal symptoms.

Talk to a doctor at once if you take too much of this medicine, even if you feel well. This is because too much paracetamol can cause delayed, serious liver damage. If your symptoms continue or your headache becomes persistent, see your doctor.

4. Possible side effects

Some people may have side effects when using this medicine. If you have any unwanted side effects you should seek advice from your doctor, pharmacist or other healthcare professional.

When using this product you may experience:

- Constipation

Stop taking this medicine and tell your doctor immediately if you experience:

- Severe abdominal pain, dry mouth, nausea and vomiting if you have recently had your gall bladder removed
- Nervousness, dizziness, irritability, drowsiness or restlessness
- Allergic reactions which may be severe such as skin rash and itching
- Skin rash or peeling or mouth ulcers, itching or sweating

! Breathing problems. These are more likely if you have experienced them before when taking other painkillers such as ibuprofen and aspirin.

- Unexplained bruising or bleeding
- Nausea, upset stomach, sudden weight loss, loss of appetite and yellowing of the eyes and skin
- Difficulty urinating.

Reporting of side effects: If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard

By reporting side effects you can help provide more information on the safety of this medicine.

! How do I know if I am addicted?

If you take this medicine according to the instructions on the pack, it is unlikely you will become addicted to the medicine. However, if the following apply to you it is important that you talk to your doctor:

- You need to take the medicine for longer periods of time.
- You need to take more than the recommended dose.
- When you stop taking the medicine you feel very unwell but you feel better if you start taking the medicine again.

5. How to store Solpadeine Plus Capsules

Keep out of the reach and sight of children.
Do not use this medicine after the 'EXP' date shown on the pack.
Store below 25°C.

6. Further information

Active ingredients: Each capsule contains Paracetamol 500 mg, Codeine Phosphate Hemihydrate 8 mg and Caffeine 30 mg.

Other ingredients: Maize starch, magnesium stearate, titanium dioxide (E 171), erythrosine (E 127), patent blue V (E 131), quinoline yellow (E 104), gelatin, shellac, propylene glycol (E 1520), iron oxide black (E 172) and ammonium hydroxide (E 527).

Packs of Solpadeine Plus Capsules contain 16 or 32 capsules.

The marketing authorisation holder is Omega Pharma Ltd, 32 Vauxhall Bridge Road, London, SW1V 2SA, United Kingdom and all enquiries should be sent to this address.

The manufacturer is SmithKline Beecham S.A., Carretera de Ajalvir, Km.2500, 28806 Alcalá De Henares, Madrid, Spain.

This leaflet was last revised in November 2013.

1100000010692

File title: 9034_SOLPC00940812v1_Solpadeine Plus capsules PIL v8F
 Item No.: 1100000010692 PROSP SOLPADEINA CAPSULAS UK 1212
 Last edit: 22nd November 2013
 Brand: Solpadeine
 Variant: Plus Capsules
 Component: Leaflet
 Market/s: UK
 Dimensions: 150 x 100mm (Drw No. L613D005)
 Printer: Santiago Gonzalez
 Substrate: White Paper
 EAN Code: N/A
 PharmaCode: 832
 Replacement number: 1100000009245
 Manufacturing Site: Alcalá
 No. of colour: 1

Process Black

81 Rivington Street
 Shoreditch
 London EC2A 3AY
 UK
 awk contact:
 Phil Hinchley
 phil@oneagency.co
 +44 (0)7710 521027
 +44 (0)203 008 6041

Regulatory Approval:

Sign:..... Date:.....

Name (please print):.....

Contact email:.....

Marketing Approval:

Sign:.....

Name (please print):.....

Contact email:.....

